

# CIDETRAK<sup>®</sup> OFM-L

**A MATING DISRUPTANT FORMULATION FOR:**  
**ORIENTAL FRUIT MOTH** (*Grapholita molesta*)  
**MACADAMIA NUT BORER** (*Cryptophlebia ambrodelta*)  
**KOA SEEDWORM** (*Cryptophlebia illepidia*)

## ACTIVE INGREDIENTS:

Z-8-Dodecen-1-yl Acetate\* .....4.65%

E-8-Dodecen-1-yl Acetate\* .....0.30%

Z-8-Dodecen-1-ol\* .....0.05%

OTHER INGREDIENTS: .....95.00%

TOTAL: .....100.00%

\*250 mgs of pheromone/dispenser

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

See additional precautionary statements on back label

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**Caution:** Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash before reuse. Wear chemical resistant gloves.

## FIRST AID STATEMENT

<b>If on skin or clothing</b>	<ul style="list-style-type: none"><li>• Remove contaminated clothing.</li><li>• Rinse skin immediately for 15-20 minutes with water.</li><li>• Contact a poison center or doctor for advice on treatment.</li></ul>
<b>If in eyes</b>	<ul style="list-style-type: none"><li>• Hold eyes open and rinse slowly and gently for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing eyes.</li><li>• Contact a poison center or doctor for advice on treatment.</li></ul>
<b>Hot line Number</b>	<ul style="list-style-type: none"><li>• When calling a poison control center or doctor, or when seeking treatment, have the product container or label with you. For additional information, you may also contact Trece Inc. at 1-866-785-1313.</li></ul>

## NET CONTENTS:

**Dispensers per package: 360 units**

*EPA Registration No. 51934-8*

*EPA Establishment No. 51934-OK-1*

## ENVIRONMENTAL HAZARDS

**For terrestrial uses:** Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of product, wash water, or when cleaning equipment.



Mailing Address: P.O. Box 129 • Adair, Oklahoma 74330  
Shipping Address: 7569 Hwy 28 W. • Adair, Oklahoma 74330  
Phone: (918) 785-3061 • Fax: (918) 785-3063

PHEROCON<sup>®</sup> and CIDETRAK<sup>®</sup> are registered Trademarks of  
TRECE Incorporated, Adair, OK USA ©2005 TRECE, Adair, OK USA



# CIDETRAK® OFM-L

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### READ ENTIRE LABEL BEFORE USING

CIDETRAK® OFM-L disrupts the mating communication between adult male and female moths, reduces fertile egg laying, suppresses subsequent larval infestation, and reduces the need for applying insecticides. For maximum effectiveness, growers remove and destroy infested fruit from trees and the ground. Growers maintain accurate records of OFM adult populations monitoring and larval damage within their orchards. Consult your local extension specialist, certified crop advisor, or Trece representative for assistance in designing the best CIDETRAK® OFM-L program for your orchard.

<b>Crop</b>	Peaches, Nectarines, Apricots, Plums, Cherries, and other stone fruits, Apples, Pears, Quince, and other pome fruits; Almonds, Macadamia, and other tree fruit crops.
<b>Targeted Pest</b>	Oriental Fruit Moth ( <i>Grapholitha molesta</i> ), Macadamia nut borer ( <i>Cryptophebia ambrodelta</i> ), Koa seedworm ( <i>Cryptophebia illepada</i> ).
<b>Rate</b>	Minimum of 100 dispensers per acre (250 dispensers per hectare) or 25.0 gm. a.i. per acre per application. Maximum of 200 dispensers per acre (500 dispensers per hectare) or 50 gm a.i. per acre per application. Apply the maximum dispenser rate in orchards that are moderately to heavily infested with OFM, or any orchard adjacent to an untreated orchard. Do not exceed 150 gm a.i. or 600 dispensers per acre per year.

<b>Application</b>	Attach dispensers securely to lateral branches in upper one-third of tree canopy. Attach the dispensers from the ground or a moving trailer using a pole applicator.
<b>Timing</b>	Apply prior to moth emergence in the spring, but no later than 2-3 days after trap capture of first male moth. Monitor moth activity using PHEROCON® 1C traps and lures. Begin monitoring in early spring and continue throughout the season to assess treatment effectiveness. Depending on temperature, <b>CIDETRAK® OFM-L</b> dispensers will release pheromone for up to 150 days. In late maturing varieties with long field seasons (i.e. more than 120 days), a second application should be made prior to subsequent OFM flights. <b>NOTE:</b> <b>CIDETRAK® OFM-L</b> will not prevent crop damage from immigration of mated female moths into treated orchards. In late season treat Oriental Fruit Moth host crops, such as pears, with pheromone only when the crops are within or near blocks that have been treated with <b>CIDETRAK® OFM-L</b> .
<b>Treatment Tips</b>	<b>CIDETRAK® OFM-L</b> suppresses mating of Oriental Fruit Moth. Immigration of mated female moths from adjacent, infested orchards can reduce the level of control. Area-wide treatment of the entire host plant block is the most effective strategy. To supplement <b>CIDETRAK® OFM-L</b> , use insecticides to control high populations of Oriental Fruit Moth. Monitor all pest populations to determine timely use of insecticides. To manage immigration and high population pressures, consider: <ol style="list-style-type: none"><li>1. Treatment of external sources of infestation with <b>CIDETRAK® OFM-L</b></li><li>2. Treatment of external sources of infestation with an insecticide</li><li>3. Treatment of pheromone treated orchard with insecticide</li></ol> Consult your local extension specialist, certified crop advisor, or Trece representative for local field condition management strategies.



## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal

**PESTICIDE STORAGE:** Store unopened original package in a dry location at temperature below 40° F. To avoid contamination of food or feed items, only unopened product packages may be stored in cold storage facilities that are used for food storage.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Dispose of empty dispensers and foil envelopes in trash.

## WARRANTY AND LIMITATION OF DAMAGES

All statements concerning the use of this product apply only when used as directed as stated on this product label. Trece Inc. warrants that the product conforms to this label and under normal conditions of use according to this label, is reasonable fit for the stated label purpose. The manufacturer makes no express or implied warranties of merchantability or fitness for a purpose concerning this product, or its use, extending beyond the label. Under no circumstance will Trece Inc. be liable for damage (special, consequential or incidental) that result from the handling, storage, or use of this product which is not in compliance with the label. Read all label directions carefully.

Buyer's exclusive remedy, should this product be defective, is replacement of the product, or if this is impractical, a purchase price refund.

