CIDETRAK® CM PP

A MATING DISRUPTION FORMULATION FOR:
CODLING MOTH *(Cydia pomonella)*
AND HICKORY SHUCKWORM *(Cydia caryana)*

ACTIVE INGREDIENT:	**WT. BY %**
8,10-Dodecadien-1-ol, (8E, 10E)	1.90%
OTHER INGREDIENTS:	98.10%
TOTAL:	**100.00%**

EPA Reg. No.: 51934–31

EPA Est. No.: 51934–OK–001

NET CONTENTS:
Dispensers per package: 400 dispensers
One dispenser contains: 95 mg 8,10-Dodecadien-1-ol, (8E,10E)

MANUFACTURED BY: TRÉCÉ INCORPORATED,
7569 Hwy 28 W, Adair, Oklahoma 74330

TRÉCÉ® INCORPORATED

Mailing Address: P.O. Box 129 • Adair, Oklahoma 74330
Shipping Address: 7569 Hwy 28 W. • Adair, Oklahoma 74330
Phone: (918) 785-3061 • Fax: (918) 785-3063

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF INHALED	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBERS:	Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at **1-800-858-7378** Mon.-Fri., 8:00 a.m. to 12:00 p.m. pacific time or your poison control center at **1-800-222-1222**. For additional information, you may also contact Trécé Inc. at **1-866-785-1313**.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes or clothing. Harmful if absorbed through the skin. Causes moderate eye and skin irritation. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:
- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of this product.

SAMPLE

CIDETRAK® CM PP

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE USING

CIDETRAK® CM PP is a dispenser system that contains behavior modifying pheromone in a polymeric matrix. It disrupts the mating communication between adult Codling Moth.

For maximum effectiveness, growers should remove and destroy infested fruit from trees and the ground. Maintain accurate records of CM adult population monitoring and larval damage within their orchards. Consult your local extension specialist, certified crop advisor, or Trécé representative for assistance in designing the best **CIDETRAK® CM PP** program for your orchard.

Crop	Apples, pears, quince and other pome fruit; walnut, pecan and other tree nuts; peaches, nectarines, plums, prunes and other stone fruits; Also, any other crop where CM is a pest.
Targeted Pest	Codling Moth (Cydia pomonella), Hickory Shuckworm (Cydia caryana).
Use Rate	**CIDETRAK® CM PP:** Depending on the population abundance, use a minimum of 200 dispensers per acre (500 dispensers per hectare) to a maximum of 400 dispensers per acre (1,000 dispensers per hectare). For example, apply 400 dispensers per acre (1,000 dispensers per hectare) for moderate to high populations. Apply the maximum dispenser rate to all orchard borders including orchards that are adjacent to an untreated or otherwise at-risk orchard heavily infested with CM. **DO NOT** exceed 150 g a.i. per acre per year.
Application	Attach dispensers securely to lateral branches in the upper 1/5th of the canopy. Distribute dispensers in a grid pattern within the orchard except on borders. Apply/attach the dispensers from the ground using a pole applicator or from a moving trailer.
Monitoring/ Timing	Apply prior to CM emergence in the spring. Monitor CM activity using PHEROCON® traps and PHEROCON® lures. Monitor throughout the season to assess treatment effectiveness. Note - **CIDETRAK® CM PP** will not prevent crop damage from immigration of CM into treated orchards. Nearby (500 yards) untreated CM hosts, such as apples, walnuts, pears, quince, crabapple, plum and any other crop where CM is a pest can be an additional source of CM. Treat with **CIDETRAK® CM PP** only when all host crops within or near treated blocks have been adequately treated for these pests.
Treatment Tips	**CIDETRAK® CM PP** suppresses mating of CM in targeted orchards. Immigration of CM from adjacent, infested orchards can reduce the level of control. Area-wide treatment of the entire host plant block is the most effective strategy. Use insecticides to supplement **CIDETRAK® CM PP** to control populations. Monitor all pest populations to determine timely use of insecticides. Consider the following to manage immigration and high population pressures: 1. Treatment of external sources of infestation with **CIDETRAK® CM PP** 2. Treatment of external sources of infestation with an insecticide 3. Treatment of pheromone treated external orchards with an insecticide Consult your local extension specialist, certified crop advisor, or Trécé representative for local field condition management strategies.

TRÉCÉ® INCORPORATED

SAMPLE

STORAGE AND DISPOSAL

DO NOT contaminate food or feed by storage or disposal of unused or used dispensers.

PESTICIDE STORAGE: Store unopened original package in a dry location at a temperature below 40°F. Store in a cool, dry place, out of direct sunlight, and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in the original container and out of the reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable container: DO NOT reuse or refill this container. Dispose of empty containers and foil envelopes in trash.

WARRANTY AND LIMITATION OF DAMAGES

To the extent consistent with applicable law, all statements concerning the use of this product apply only when used as directed as stated on this product label. Trécé®, Inc. warrants that the product conforms to this label and under normal conditions of use according to this label, is reasonably fit for the stated label purpose. The manufacturer makes no express or implied warranties of merchantability or fitness for a purpose concerning this product, or its use, extending beyond the label. Under no circumstance will Trécé®, Inc. be liable for damage (special, consequential, or incidental) that result from the handling, storage, or use of this product which is not in compliance with the label. Read all label directions carefully.

Buyer's exclusive remedy, should this product be defective, is replacement of the product, or if this is impractical, a purchase price refund.

© 2025, Trécé Inc., Adair, OK USA · ® is a registered trademark and TM is a trademark of Trécé Inc., Adair, OK 74330 USA