PHEROCON®

BROWN MARMORATED STINK BUG

PHEROCON BMSB DUAL™ LURES also attract:

Green Stink Bug (GSB), *Acrosternum hilare*,
Brown-Winged Green Bug, *Plautia stali*, and other Stink Bugs

Brown Marmorated Stink Bug (BMSB), *Halyomorpha halys*

Green Stink Bug (GSB), *Acrosternum hilare*

Brown-Winged Green Bug, *Plautia stali*

Brown Marmorated Stink Bug (BMSB)
Pear damage

Brown Marmorated Stink Bug (BMSB)
Raspberries damage

Stink bugs and other plant bugs may differ in color and size, however, all have the same overall shield-shaped body.

TRÉCÉ® INCORPORATED
INSECT PHEROMONE & KAIROMONE SYSTEMS
Your Edge - And Ours - Is Knowledge.

PHEROCON® BROWN MARMORATED STINK BUG

COMPLETE MONITORING SYSTEMS FOR EARLIER DETECTION OF ADULT & NYMPHAL STINK BUGS

Brown Marmorated Stink Bug (BMSB), *Halyomorpha halys*, **Green Stink Bug (GSB)**, *Acrosternum hilare*, **Brown-Winged Green Bug**, *Plautia stali*, and other Stink Bugs



STKY™ Dual Panel Adhesive Trap attached to stake with twist ties or staples



PHEROCON® BMSB DUAL Lures



PHEROCON BMSB STKY Liner



PHEROCON BMSB STKY CANOPY™ Trap can be attached to elbow stake - experimental







Guidelines For Use:

PHEROCON Brown Marmorated Stink Bug:

Placement Time:

Adults can be detected throughout the active growing season.

Placement Pattern:

- *At least 2 traps per orchard or vineyard: 1 trap per susceptible border. Do not place on tree or vine.*
- *Ensure that the trap has a reasonable surface contact with the stake by securing it horizontally along the stake.*

Row Crops:

- *At least 2 traps per planting; 1 trap per susceptible border.*
- *Only place traps along risky borders. Do not place traps in the middle of the block or where traps are touching the crop.*
- *Place traps closer to wooded habitats rather than up against the crop.*
- *If the crop is completely isolated, place traps at the edge of the crop.*
- *Traps can be placed on stakes approximately 2-3 feet above the ground.*

Trap Maintenance:

- *Check traps weekly.*
- *Change traps/liners every 3-4 weeks or whenever they are dirty. Change lures every 14 weeks.*

Lure Storage:

- *Keep unopened factory-sealed packages in a cool place (< 75˚F or 24˚C maximum) for short periods until used. Refrigerate or freeze unused packages.*
- *Refrigerate or freeze unopened packages to carry over for annual storage.*

Contact your supplier, local extension agent or farm advisor for more details!

Visit our website or call toll-free: 1-866-785-1313

www.trece.com

TRÉCÉ® INCORPORATED
INSECT PHEROMONE & KAIROMONE SYSTEMS
Your Edge - And Ours - Is Knowledge.

© 2025, Trécé Inc., Adair, OK USA • ® is a registered trademark and TM is a trademark of Trécé Inc., Adair, OK USA